## IMPACT OF HIGH SPEED CIVIL TRANSPORTS ON STRATOSPHERIC OZONE: A 2-D MODEL INVESTIGATION

<u>Douglas E. Kinnison</u>, Peter S. Connell, Keith E. Grant, and Douglas A. Rotman Lawrence Livermore National Laboratory Livermore CA (kinnison1@llnl.gov)

## **Abstract**

This study investigates the effect on stratospheric ozone from a fleet of proposed High Speed Civil Transports (HSCTs). The new LLNL 2-D operator-split chemical-radiative-transport model of the troposphere and stratosphere is used for this HSCT investigation. This model is integrated in a diurnal manner, using an implicit numerical solver (SMVGEARII, Jacobson, 1995). Therefore, rate coefficients derived during a model integration are not modified by any sort of diurnal average factor (unlike the previous LLNL 2-D model architecture). This model, like previous versions of the LLNL 2-D model, also does not make any assumptions on lumping of chemical species into families. Comparisons to model-derived HSCT assessment of ozone change are made, both to the previous LLNL 2-D model and to other models from the international assessment modeling community (Stolarski, et al., 1995). In addition, the new LLNL 2-D model chemical solution approach is evaluated against data taken from a recent ER2 aircraft campaign. Here, aircraft measurements of simultaneous trace constituents distributions during the Stratospheric Photochemistry, Aerosol and Dynamics Expedition (SPADE), allow one to determine the odd-oxygen loss partitioning among chemical families (Wennberg et al., 1994). Very good agreement for odd-oxygen loss partitioning is found between the model and the SPADE data. In addition, the sensitivity to the NO<sub>x</sub> emission index and sulfate surface area density is also explored.

Stolarski, R.et al., 1995 Scientific Assessment of the Atmospheric Effects of Stratospheric Aircraft, NASA Ref. Publ. 1381, November 1995.

Jacobson, M. Z., Computation of global photochemistry with SMVGEARII, *Atm. Env.*, 29, 2541-2546, 1995.

Wennberg, P.O., et al., Removal of Stratospheric O<sub>3</sub> by Radicals: In Situ Measurements of OH, HO<sub>2</sub>, NO, NO<sub>2</sub>, ClO, and BrO, *Science*, 266, 398-404, 1994.

\*This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.